

U. PORTO

FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2017/2018

Catarina Metelo Coimbra dos Santos Ferreira
Placenta Artificial: Avanços Recentes e
Potenciais Aplicações Clínicas
Artificial Placenta: Recent Advances
and Potential Clinical Applications

março, 2018

FMUP

Catarina Metelo Coimbra dos Santos Ferreira
Placenta Artificial: Avanços Recentes e
Potenciais Aplicações Clínicas
Artificial Placenta: Recent Advances
and Potential Clinical Applications

Mestrado Integrado em Medicina

Área: Fisiologia

Tipologia: Monografia

**Trabalho efetuado sob a Orientação de:
Professor Doutor Roberto Roncon-Albuquerque**

**Trabalho organizado de acordo com as normas da revista:
Pediatric Pulmonology**

março, 2018

Eu, **CATARINA METELO COIMBRA DOS SANTOS FERREIRA**, abaixo assinado, nº mecanográfico **201200152**, estudante do 6.º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 20/03/2018

Assinatura conforme cartão de identificação:

Catarina Metelo Coimbra dos Santos Ferreira

NOME

CATARINA METELO COIMBRA DOS SANTOS FERREIRA

NÚMERO DE ESTUDANTE

201200152

E-MAIL

catarinametelocoimbra@hotmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

FISIOLOGIA

TÍTULO MONOGRAFIA

ARTIFICIAL PLACENTA: RECENT ADVANCES AND POTENTIAL CLINICAL APPLICATIONS

ORIENTADOR

PROFESSOR DOUTOR ROBERTO RONCON-ALBUQUERQUE

COORIENTADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 20/03/2019

Assinatura conforme cartão de identificação: Catarina Metelo Coimbra dos Santos Ferreira

ARTIFICIAL PLACENTA: RECENT ADVANCES AND POTENTIAL CLINICAL APPLICATIONS

Catarina Metelo-Coimbra and Roberto Roncon-Albuquerque Jr^{1,2}

1. Department of Physiology and Cardiothoracic Surgery, Faculty of Medicine of Porto, PORTUGAL

2. Department of Emergency and Intensive Care Medicine, Hospital de S. João, Porto, PORTUGAL

Abbreviated Title: Recent Advances in the Artificial Placenta.

Correspondence address:

Roberto Roncon-Albuquerque Jr, MD PhD

Department of Physiology and Cardiothoracic Surgery, Faculty of Medicine of Porto and

Department of Emergency and Intensive Care Medicine Hospital de S. João

Al. Prof. Hernâni Monteiro, Porto 4200-319, PORTUGAL

Tel.: +351 916 454 074

Fax: +351 225 025 766

E-mail: rra_jr@yahoo.com

ABSTRACT

Lung immaturity remains a major cause of morbidity and mortality in extremely premature infants. Positive-pressure mechanical ventilation, the method of choice for respiratory support in premature infants, frequently promotes by itself lung injury and a negative impact in the circulatory function. Extracorporeal lung support has been proposed for more than 50 years as a potential alternative to mechanical ventilation in the treatment of severe respiratory failure of extremely premature infants. Recent advances in this field included the development of miniaturized centrifugal pumps and polymethylpentene oxygenators, as well as the successful use of pump-assisted venovenous extracorporeal gas exchange systems in experimental artificial placenta models. This review, which includes studies published from 1958 to 2015, presents an update on the artificial placenta concept and its potential clinical applications. Special focus will be devoted to the milestones achieved so far and to the limitations that must be overcome before its clinical application. Notwithstanding, the artificial placenta stands as a promising alternative to mechanical ventilation in extremely premature infants.

Keywords: Artificial placenta; bronchopulmonary dysplasia; extracorporeal membrane oxygenation; extremely premature infant; respiratory distress syndrome.

INTRODUCTION

According to the World Health Organization, 15 million premature infants are born every year, with almost 1 million deaths directly attributed to prematurity¹. Although all premature infants are at risk for complications, extremely premature (EPT) infants, born at or before 28 weeks of gestation, suffer the greatest morbidity and mortality¹. Respiratory distress syndrome and bronchopulmonary dysplasia remain a major cause of morbidity and mortality in EPT infants². This relates with pulmonary immaturity, given that EPT infants are born during the canalicular period of lung development, characterized by capillarization and pulmonary acini morphogenesis, with insufficient surfactant production³.

Positive-pressure mechanical ventilation with high oxygen concentrations remains the method of choice to provide lung support in preterm infants with severe respiratory failure⁴. However, mechanical ventilation in preterm infants promotes, by itself, lung injury that negatively impacts survival. The known direct mechanisms of ventilator-induced lung injury are barotrauma, volutrauma and atelectrauma⁵. These mechanisms promote biotrauma with capillary endothelium, alveolar epithelium, and basal membrane damage, which results in fluid, protein and blood extravasation into the airways, alveoli, and pulmonary interstitium, with consequent surfactant inhibition and activation of local and systemic inflammatory responses⁶. Mechanical ventilation also adversely affects the circulatory function of preterm infants, reducing pulmonary blood flow and left ventricular output^{7,8}. In fact, positive-pressure mechanical ventilation decreases the alveolar/capillary transmural pressure gradient, causing compression of the intra-alveolar capillaries, which increases pulmonary vascular resistance, therefore decreasing pulmonary blood flow⁹. Further increases in airway pressure and pulmonary vascular resistance may sustain pulmonary arterial pressures above systemic arterial

pressures, potentiating continued right-to-left shunt through the ductus arteriosus. The effect of positive-pressure ventilation is not limited to the pulmonary vasculature, with direct compressive effects observed on the newborn heart, resulting in reduced cardiac performance and ventricular output¹⁰. This could be particularly relevant in the immature myocardium of the preterm heart that presents low contractility with an inability to cope with increasing afterload in the days after birth¹¹.

In light of these limitations, extracorporeal lung support has been proposed more than 50 years ago as a potential alternative to mechanical ventilation in the treatment of severe respiratory failure of EPT infants¹². Its main benefits reside in the fact that, by bypassing the lungs completely, the AP avoids potential barotrauma resulting from mechanical ventilation. However, by then, artificial organ technology was still in its infancy and the understanding of EPT pathophysiology was very limited. More recently, important developments in extracorporeal lung assist technology were observed, fostering a renewed interest in artificial placenta (AP) research.

In this review, an update on the AP concept and its potential applications is presented. Special focus will be devoted to the milestones achieved so far and to the major limitations that must be overcome before its clinical application.

METHODS

Eligible studies were identified by an electronic search of PubMed and Scopus, involving studies published from 1958 to 2015. The sensitive search strategy combined the following keywords: *artificial placenta*; *bronchopulmonary dysplasia*; *extracorporeal lung assist*; *extracorporeal membrane oxygenation*; *extremely premature infants*; *polymethylpentene oxygenator*; and *respiratory distress syndrome*. All articles and cross-referenced studies from retrieved articles were screened for pertinent information and reviewed by both Authors.

Inclusion criteria consisted in experimental and systematic review articles, published as original studies, with available abstract. Publications not written in English or not related to the neonatal period were excluded.

ARTIFICIAL PLACENTA: THE CONCEPT

The terminology describing extracorporeal life support (ECLS) in premature infants is divided in three categories, established according to the gestational age¹³: *i*) neonatal ECMO: for infants of at least 34 weeks of gestation (moderate to late preterm newborns¹); this technique is successfully used in clinical practice for more than 30 years; *ii*) preemie ECMO: for premature infants between 29 and 33 weeks of gestation (very preterm newborns); although technically feasible, reduced survival and increased rates of intra-ventricular hemorrhage have been reported¹⁴; and *iii*) artificial placenta: for EPT infants, born before 28 weeks; the AP currently remains under experimental research.

Therefore, the AP concept consists in extracorporeal membrane gas exchange (blood oxygenation and extracorporeal CO₂ removal) for EPT infants with immature pulmonary system and severe respiratory failure, as a bridge to the development of native lung function⁸. The AP does not include provision of other placental functions such as nutrient or metabolic product exchange.

AP models are generally defined by the following characteristics^{8,13,15,16}: *i*) extracorporeal lung support with preservation of fetal circulation: either the umbilical artery (pumpless arterio-venous ECLS, AV-ECLS) or a central vein (pump-assisted veno-venous ECLS, VV-ECLS) are used for blood outflow from the patient to the extracorporeal circuit; the umbilical vein is always used for inflow to the patient from the extracorporeal circuit¹⁷ (Fig. 1); *ii*) low partial pressure of oxygen, given that oxygen-binding capacity of fetal hemoglobin and hematocrit are increased; *iii*) absence of positive-pressure mechanical ventilation; *iv*) simulated fetal breathing with fluid-filled lungs; *v*) biocompatibility of the extracorporeal circuit inner surfaces in direct

contact with blood; high levels of systemic anticoagulation associated with an unacceptable risk of intracranial hemorrhage in EPT infants¹⁸.

Pump-assisted VV-ECLS systems presents several advantages when compared to pumpless AV-ECLS¹⁷: *i*) arterial vessel cannulation is not required; cannulation of umbilical arteries in EPT infants is technically challenging and frequently complicates with vessel spasm; *ii*) an external blood reservoir is not required given that the EPT infant's own venous system is used as blood reservoir; *iii*) it operates in parallel with systemic circulation, therefore not increasing afterload of the fetal heart; differently, pumpless AV-ECLS operates in series with systemic circulation, increasing the fetal heart workload that can complicate with high-output heart failure.

Membrane lung permeability to O₂ and CO₂ is a critical factor influencing AP performance. Silicone rubber membranes were initially used, composed of a permeable non-porous polymeric material with loosely packed polymeric chains¹⁹. Another type of oxygenator is the hollow-fiber membrane, which has woven capillary tubes composed of microporous polypropylene²⁰. Despite higher gas permeability, polypropylene membranes are less frequently used given the high plasma leakage risk with increased extracorporeal circuit blood pressures²⁰. Moreover, silicone rubber membranes present higher biocompatibility when compared with polypropylene fibers for long-term ECLS²¹.

Polymethylpentene (PMP) oxygenators have been used in clinical practice for a decade²². PMP fibers have an asymmetric pore structure with a very thin dense outer skin that allows gas transference while suppressing the direct contact of blood and gas across micropores²³. These features enhance durability and greatly reduce plasma leakage²³. Moreover, PMP oxygenators present more efficient priming, reduced hemodynamic resistance and better preservation of coagulation proteins²¹.

EARLY MILESTONES

Following the development and implementation of the heart-lung machine, it was soon recognized that this concept could be similarly applied to the treatment of severe respiratory failure of the premature infant²⁴. In 1958, Westin et al. prolonged the life of previable human fetuses by cannulating the umbilical vessels and circulating the fetal blood through a rotating oxygenator²⁵. When injecting regular doses of glucose solution, the fetal heart continued to beat for a period up to 12 hours²⁵. The fetuses were maintained in a warm artificial amniotic fluid bath²⁵.

Callaghan et al. were the first to develop the AP concept in animal experiments, back in 1961. A pump-assisted VV-ECLS circuit with a rotating disc oxygenator was used in eight sheep²⁶. Blood outflow from the animal to the extracorporeal circuit was performed via both femoral and jugular veins, while inflow to the animal from the extracorporeal circuit was made either through the right atrium or the right ventricle²⁶. In 1962, a period of up to 2.5 hours survival of mongrel dogs using this procedure was reported²⁷.

These achievements have been overcome by Lawn and McCance²⁸, who conceived a pumpless AV-ECLS circuit with a dialyzer that was tested in previable pig fetuses. Blood drained from the umbilical arteries circulated through the oxygenator and then through cellophane tubing immersed in a suitable rinse. Blood returned to the umbilical vein without requiring external pump assistance²⁸. A similar perfusion apparatus was constructed in 1964 by Alexander et al., but the dialyzer was excluded²⁹. It was concluded from their experiments that a perfusion system with constant volume would be necessary, due to changes in venous pressure²⁹.

Meanwhile, SenGupta et al. described a portable, self-contained and self-powered AP consisting of a flexible, inert silicone elastomer membrane oxygenator and a pump³⁰. Their first experiment had eleven out of sixteen survivors during up to 5 hours of connection to the AP³⁰. Two years later, there were sixteen survivors out of twenty dogs, and most perfusions took more than 2 hours. Though satisfactory oxygenation was obtained, the short period of survival was considered unsafe³¹.

Significant progress was achieved in 1969 by Zapol et al.³², when a premature lamb fetus was totally sustained by extracorporeal perfusion using a silicone-membrane blood oxygenator³². The animal received parenteral nutritional support and remained metabolically stable for up to 55 hours. A study with ten lamb fetuses using angiocardiographic techniques was presented the following year³³. Zapol et al. also described the modulation of ductus arteriosus and pulmonary blood flow by blood oxygen tension in their AP model³⁴.

Efforts on the development of an AP were almost entirely abandoned by 1979, when a completely different approach to treat severe respiratory failure in premature infants was implemented: positive-pressure mechanical ventilation^{35,36}. This led to a dramatic improvement of premature infants' survival, although many problems remained³⁷. Compared with positive-pressure mechanical ventilation, the AP was at that time too complex and unsafe for clinical use leading to a gap of research in the next decade.

In 1987, Kuwabara et al.³⁸ developed a novel AP system. They compared two types of circuits, with and without a blood reservoir, using goat fetuses. In the first group, the duration of incubation was increased to 165 hours, in contrast to the 8 hours achieved by the control group³⁸. The oxygenator was made of silicone, and blood was drained from the umbilical arteries and returned to the umbilical vein³⁸. This was the

first report on animal experiments of successful long-term (up to 7 days) AP support. Several improvements were made to this novel AP system, including alterations in fetal catheterization and addition of a dialyzing system to the extracorporeal circuit, which allowed the survival of goat fetuses up to 236 hours³⁹.

In 1993, continuing this work, Unno et al. tried a new protocol to study the influence of body movement on goat fetuses' survival⁴⁰. It was demonstrated that AV-ECLS with umbilical blood access could support premature goat fetuses for up to 3 weeks⁴⁰. The following studies focused on fetal hemodynamics, such as goat fetal ductal blood velocity through Doppler echocardiography⁴¹ and the effect of prostaglandin E1⁴². This last experiment suggested that the administration of prostaglandin E1 prevented the constriction of ductus arteriosus, a phenomenon that was found to disturb fetal circulation to the AP⁴³. This research group also compared four different methods to control blood flow in an AV-ECLS circuit, concluding that the control of extracorporeal circulation flow by altering the circuit resistance was one of the main contributing factors to the success of long-term incubation⁴⁴.

In 1998, Sakata et al. reported the successful use of a centrifugal pump, which allowed higher extracorporeal flow rates⁴⁵. This group used polyolefin hollow fiber membrane oxygenators, which contributed to low circuit resistance⁴⁵. In the same year, Yasufuku et al. refined the AP concept by performing upper tracheal ligation, which maintained lung expansion and protected from meconium aspiration^{46,47}.

RECENT MILESTONES

In 2012 Gray et al.⁴⁸, from the University of Michigan ECLS Laboratory, hypothesized that a pump-assisted VV-ECLS circuit would preserve systemic fetal circulation while providing adequate extracorporeal gas exchange⁴⁸. The right jugular vein was cannulated for outflow from the animal to the extracorporeal circuit, whereas an umbilical vein was used for blood inflow to the animal from the extracorporeal circuit⁴⁸. Blood cavitation was reduced, since blood was passively drained from the right atrium⁴⁸. A miniaturized polypropylene hollow fiber oxygenator was used. The experiment was successful, being the first report of a 24-hour survival of five lamb fetuses using a pump-assisted VV-ECLS circuit. Continuing this work, seven lambs were incubated on a dry heated waterbed and maintained on VV-ECLS for up to 70 hours⁴⁹. This AP model was able to provide hemodynamic stability and efficient extracorporeal gas exchange, with preservation of cerebral perfusion for an extended period of time⁴⁹. No signs of gross or microscopic intra-ventricular hemorrhage were found despite systemic anticoagulation with heparin⁴⁹.

In 2015, Bryner et al. compared a pump-assisted VV-ECLS system with positive-pressure mechanical ventilation in EPT lamb fetuses¹⁶. Four lambs were successfully supported for 1 week using a polypropylene oxygenator and a rotary pump¹⁶. Differently, lambs treated with positive-pressure mechanical ventilation survived on average less than 4 hours, despite the use of exogenous surfactant and steroids¹⁶. No evidence of intracranial hemorrhage was observed. The main issues faced by researchers were directly related to cannulation, with one case of pericardial tamponade and arrhythmias¹⁶.

Besides the improvements in extracorporeal circuit configuration, important advances were achieved in oxygenator technology. Arens et al.⁵⁰ developed a

miniaturized oxygenator to be used in AP models. This research group catheterized lambs through two umbilical arteries and two umbilical veins, and the fetuses were kept in a warming bed. Their oxygenator, NeonatOx, was placed as close as possible to the lambs, allowing short tubing and low circuit resistance⁵⁰. Furthermore, the oxygenator was miniaturized to a priming volume of only 20 ml. This reduced device surface area, decreasing thrombogenesis and inflammation⁵⁰. NeonatOx allowed successful extracorporeal gas exchange for 6 hours in six out of seven animals⁵¹. One limitation related to the durability of umbilical vascular accesses, since artificial amniotic medium submersion was not performed⁵¹.

Meanwhile, Canadian investigators designed a microfluidic oxygenator with efficient gas exchange²⁰. Four different gas permeable membranes were tested using human blood. The porous polydimethylsiloxane membrane had the highest gas exchange rate²⁰. Recently, further improvements were performed to this oxygenator⁵², with the novel device being modifiable according to the EPT infants' body weight. This microfluidic oxygenator was tested in piglets during 4 hours⁵².

FUTURE DIRECTIONS

Although much progress has been made in the AP field and despite the different models studied so far (Table 1 and Table 2), several limitations still preclude its clinical application.

Regarding the extracorporeal circuit itself, pump-assisted VV-ECLS models have shown many advantages over pumpless AV-ECLS circuits. Although its simplicity is appealing, the use of pumpless AV-ECLS in EPT infants seems technically impracticable due to the small size and tortuosity of the umbilical arteries, as well as to the need of prolonged extracorporeal lung support and hemodynamic stability.

Concerning anticoagulation, the development of novel biomaterials will presumably improve surface biocompatibility of the extracorporeal circuit, reducing (or even eliminating) the need for systemic anticoagulation, importantly decreasing intracranial hemorrhage risk¹⁸. In this regard, research is underway towards the development of non-thrombogenic surface extracorporeal circuit coating^{53,54}.

Artificial placenta miniaturization will also be improved, decreasing extracorporeal surface area and circuit resistance⁵¹. The use of PMP oxygenators, which present high durability and reduced plasma leakage, is a predictable next step, given its successful use in adult ECMO²³.

Further studies are required to show that the lung is protected and continues to mature during AP support. This implies that lung development between the stages of birth, AP support and progression to air breathing needs to be demonstrated and documented.

Concerning the brain, studies need to show that there is adequate brain perfusion and that this organ is protected without bleeding or white matter injury during AP

support. This is essential given that, regarding neurological complications, the majority of sequelae appear to be related to hypoxemia and hemodynamic instability that occurs before the onset of ECLS⁵⁵.

The impact of the AP in the cardiovascular, gastrointestinal and renal systems also deserves further investigation. Cardiovascular stability during pump-assisted VV-ECLS in EPT infants also needs to be confirmed, before establishing this configuration as the preferred AP circuit.

The ability of EPT infants to wean from AP support without major lung sequelae is a central issue that needs further demonstration. This will impact on clinical criteria for AP use, which remain to be established.

CONCLUSION

Lung immaturity still associates with high morbidity and mortality in EPT infants. Extracorporeal lung support has been proposed for more than 50 years as a potential treatment of severe respiratory failure of EPT infants. Recent progresses in extracorporeal circuit biotechnology renewed the interest in experimental and clinical AP research. Notwithstanding the several challenges remaining, the AP remains an attractive potential alternative for EPT infants failing positive-pressure mechanical ventilation. The successful application of the AP into clinical practice would definitely be a milestone in neonatal medicine.

REFERENCES

1. Howson CP, Kinney MV, McDougall L, Lawn JE, Born Too Soon Preterm Birth Action G. 2013. Born too soon: Preterm birth matters. *Reprod Health*. 10 Suppl 1:S1.
2. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, Hale EC, Newman NS, Schibler K, Carlo WA et al. 2010. Neonatal outcomes of extremely preterm infants from the nichd neonatal research network. *Pediatrics*. 126(3):443-456.
3. Davis RP, Mychaliska GB. 2013. Neonatal pulmonary physiology. *Semin Pediatr Surg*. 22(4):179-184.
4. Committee on F, Newborn, American Academy of P. 2014. Respiratory support in preterm infants at birth. *Pediatrics*. 133(1):171-174.
5. Carvalho CG, Silveira RC, Procianoy RS. 2013. Ventilator-induced lung injury in preterm infants. *Rev Bras Ter Intensiva*. 25(4):319-326.
6. Dreyfuss D, Saumon G. 1998. Ventilator-induced lung injury: Lessons from experimental studies. *Am J Respir Crit Care Med*. 157(1):294-323.
7. Kluckow M, Evans N. 1996. Relationship between blood pressure and cardiac output in preterm infants requiring mechanical ventilation. *The Journal of pediatrics*. 129(4):506-512.

8. Rochow N, Chan EC, Wu WI, Selvaganapathy PR, Fusch G, Berry L, Brash J, Chan AK, Fusch C. 2013. Artificial placenta--lung assist devices for term and preterm newborns with respiratory failure. *Int J Artif Organs*. 36(6):377-391.
9. Lang JA, Pearson JT, Binder-Heschl C, Wallace MJ, Siew ML, Kitchen MJ, Te Pas AB, Fouras A, Lewis RA, Polglase GR et al. 2015. Increase in pulmonary blood flow at birth; role of oxygen and lung aeration. *J Physiol*.
10. Biondi JW, Schulman DS, Soufer R, Matthay RA, Hines RL, Kay HR, Barash PG. 1988. The effect of incremental positive end-expiratory pressure on right ventricular hemodynamics and ejection fraction. *Anesth Analg*. 67(2):144-151.
11. Takahashi Y, Harada K, Kishkurno S, Arai H, Ishida A, Takada G. 1997. Postnatal left ventricular contractility in very low birth weight infants. *Pediatr Cardiol*. 18(2):112-117.
12. Callaghan JC, Maynes EA, Hug HR. 1965. Studies on lambs of the development of an artificial placenta. Review of nine long-term survivors of extracorporeal circulation maintained in a fluid medium. *Canadian journal of surgery Journal canadien de chirurgie*. 8:208-213.
13. Davis RP, Bryner B, Mychaliska GB. 2014. A paradigm shift in the treatment of extreme prematurity: The artificial placenta. *Curr Opin Pediatr*. 26(3):370-376.
14. Kim A MK, Rana A, Drongowski R, Bartlett R, Hirschl R, Mychaliska G. 2009. Pushing the boundaries of ecls: Outcomes in < 34 week ega neonates. Paper

presented at: American Pediatric Surgical Association Fortieth Annual Meeting.

American Pediatric Surgical Association; Puerto Rico

15. Bryner BS, Mychaliska GB. 2014. Ecls for preemies: The artificial placenta. *Semin Perinatol.* 38(2):122-129.
16. Bryner B, Gray B, Perkins E, Davis R, Hoffman H, Barks J, Owens G, Bocks M, Rojas-Pena A, Hirschl R et al. 2015. An extracorporeal artificial placenta supports extremely premature lambs for 1 week. *Journal of pediatric surgery.* 50(1):44-49.
17. Gray BW, Shaffer AW, Mychaliska GB. 2012. Advances in neonatal extracorporeal support: The role of extracorporeal membrane oxygenation and the artificial placenta. *Clin Perinatol.* 39(2):311-329.
18. Awad JA, Cloutier R, Fournier L, Major D, Martin L, Masson M, Guidoin R. 1995. Pumpless respiratory assistance using a membrane oxygenator as an artificial placenta: A preliminary study in newborn and preterm lambs. *J Invest Surg.* 8(1):21-30.
19. Robb WL. 1968. Thin silicone membranes--their permeation properties and some applications. *Ann N Y Acad Sci.* 146(1):119-137.
20. Wu WI, Rochow N, Chan E, Fusch G, Manan A, Nagpal D, Selvaganapathy PR, Fusch C. 2013. Lung assist device: Development of microfluidic oxygenators for preterm infants with respiratory failure. *Lab Chip.* 13(13):2641-2650.

21. Khoshbin E, Westrope C, Pooboni S, Machin D, Killer H, Peek GJ, Sosnowski AW, Firmin RK. 2005. Performance of polymethyl pentene oxygenators for neonatal extracorporeal membrane oxygenation: A comparison with silicone membrane oxygenators. *Perfusion*. 20(3):129-134.
22. Peek GJ, Killer HM, Reeves R, Sosnowski AW, Firmin RK. 2002. Early experience with a polymethyl pentene oxygenator for adult extracorporeal life support. *ASAIO J*. 48(5):480-482.
23. Toomasian JM, Schreiner RJ, Meyer DE, Schmidt ME, Hagan SE, Griffith GW, Bartlett RH, Cook KE. 2005. A polymethylpentene fiber gas exchanger for long-term extracorporeal life support. *ASAIO J*. 51(4):390-397.
24. Schoberer M, Arens J, Lohr A, Seehase M, Jellema RK, Collins JJ, Kramer BW, Schmitz-Rode T, Steinseifer U, Orlikowsky T. 2012. Fifty years of work on the artificial placenta: Milestones in the history of extracorporeal support of the premature newborn. *Artif Organs*. 36(6):512-516.
25. Westin B, Nyberg R, Enhorning G. 1958. A technique for perfusion of the previable human fetus. *Acta Paediatr*. 47(4):339-349.
26. Callaghan JC, Angeles JD. 1961. Long-term extracorporeal circulation in the development of an artificial placenta for respiratory distress of the newborn. *Surg Forum*. 12:215-217.

27. Callaghan JC, Cardozo D, Boracchia B, Aleksiuksa. 1962. Study of prepulmonary bypass in the development of an artificial placenta for prematurity and respiratory distress syndrome of the newborn. *The Journal of thoracic and cardiovascular surgery*. 44:600-607.
28. Lawn L, McCance RA. 1962. Ventures with an artificial placenta. I. Principles and preliminary results.
29. Alexander DP, Britton HG, Nixon DA. 1964. Survival of the foetal sheep at term following short periods of perfusion through the umbilical vessels. *J Physiol*. 175:113-124.
30. SenGupta A, Taylor HP, Kolff WJ. 1964. An artificial placenta designed to maintain life during cardiorespiratory distress. *Trans Am Soc Artif Intern Organs*. 10:63-65.
31. Sarin CL, SenGupta A, Taylor HP, Kolff WJ. 1966. Further development of an artificial placenta with the use of membrane oxygenator and venovenous perfusion. *Surgery*. 60(3):754-760.
32. Zapol WM, Kolobow T, Pierce JG, Bowman RL. 1969. Artificial placenta: Two days of total extrauterine support of the isolated premature lamb fetus. *Science*. 166(3905):617-618.
33. Doppman JL, Zapol W, Kolobow T, Pierce J. 1970. Angiocardiography of fetal lambs on artificial placenta. *Invest Radiol*. 5(3):181-186.

34. Zapol WM, Kolobow T, Doppman J, Pierce JE. 1971. Response of ductus arteriosus and pulmonary blood flow to blood oxygen tension in immersed lamb fetuses perfused through an artificial placenta. *The Journal of thoracic and cardiovascular surgery*. 61(6):891-903.
35. Speidel BD, Dunn PM. 1976. Use of nasal continuous positive airway pressure to treat severe recurrent apnoea in very preterm infants. *Lancet*. 2(7987):658-660.
36. Kirby RR. 1977. Intermittent mandatory ventilation in the neonate. *Crit Care Med*. 5(1):18-22.
37. Gerhardt T, Reifenberg L, Goldberg RN, Bancalari E. 1989. Pulmonary function in preterm infants whose lungs were ventilated conventionally or by high-frequency oscillation. *The Journal of pediatrics*. 115(1):121-126.
38. Kuwabara Y, Okai T, Imanishi Y, Muronosono E, Kozuma S, Takeda S, Baba K, Mizuno M. 1987. Development of extrauterine fetal incubation system using extracorporeal membrane oxygenator. *Artif Organs*. 11(3):224-227.
39. Kuwabara Y, Okai T, Kozuma S, Unno N, Akiba K, Shinozuka N, Maeda T, Mizuno M. 1989. Artificial placenta: Long-term extrauterine incubation of isolated goat fetuses. *Artif Organs*. 13(6):527-531.
40. Unno N, Kuwabara Y, Okai T, Kido K, Nakayama H, Kikuchi A, Narumiya Y, Kozuma S, Taketani Y, Tamura M. 1993. Development of an artificial placenta:

Survival of isolated goat fetuses for three weeks with umbilical arteriovenous extracorporeal membrane oxygenation. *Artif Organs*. 17(12):996-1003.

41. Sakata M, Hisano K, Okada M, Yasufuku M, Yokoyama N, Uetani Y, Noma O, Nakamura H. 1994. Experimental study of artificial placenta; doppler echocardiographic evaluation of goat fetal ductal blood velocity and waveforms in artificial placenta. *Japanese Journal of Artificial Organs*. 23(3):921-924.
42. Sakata M, Hisano K, Okada M, Yasufuku M, Yokoyama N, Uetani Y, Nakamura H. 1995. Effect of prostaglandin e1 for fetal circulation in artificial placenta. *Japanese Journal of Artificial Organs*. 24(2):604-607.
43. Sakata M, Yasufuku M, Hisano K, Okada M. 1996. Fetal circulation under the condition of artificial placenta with av-ecmo. *Japanese Journal of Artificial Organs*. 25(2):290-293.
44. Unno N, Baba K, Kozuma S, Nishina H, Okai T, Kuwabara Y, Taketani Y. 1997. An evaluation of the system to control blood flow in maintaining goat fetuses on arterio-venous extracorporeal membrane oxygenation: A novel approach to the development of an artificial placenta. *Artif Organs*. 21(12):1239-1246.
45. Sakata M, Hisano K, Okada M, Yasufuku M. 1998. A new artificial placenta with a centrifugal pump: Long-term total extrauterine support of goat fetuses. *The Journal of thoracic and cardiovascular surgery*. 115(5):1023-1031.

46. Yasufuku M, Hisano K, Sakata M, Okada M. 1998. Arterio-venous extracorporeal membrane oxygenation of fetal goat incubated in artificial amniotic fluid (artificial placenta): Influence on lung growth and maturation. *Journal of pediatric surgery*. 33(3):442-448.
47. Carmel JA, Friedman F, Adams FH. 1965. Fetal tracheal ligation and lung development. *Am J Dis Child*. 109:452-456.
48. Gray BW, El-Sabbagh A, Rojas-Pena A, Kim AC, Gadepali S, Koch KL, Capizzani TR, Bartlett RH, Mychaliska GB. 2012. Development of an artificial placenta iv: 24 hour venovenous extracorporeal life support in premature lambs. *ASAIO J*. 58(2):148-154.
49. Gray BW, El-Sabbagh A, Zakem SJ, Koch KL, Rojas-Pena A, Owens GE, Bocks ML, Rabah R, Bartlett RH, Mychaliska GB. 2013. Development of an artificial placenta v: 70 h veno-venous extracorporeal life support after ventilatory failure in premature lambs. *Journal of pediatric surgery*. 48(1):145-153.
50. Arens J, Schoberer M, Lohr A, Orlikowsky T, Seehase M, Jellema RK, Collins JJ, Kramer BW, Schmitz-Rode T, Steinseifer U. 2011. Neonatox: A pumpless extracorporeal lung support for premature neonates. *Artif Organs*. 35(11):997-1001.
51. Schoberer M, Arens J, Erben A, Ophelders D, Jellema RK, Kramer BW, Bruse JL, Brouwer P, Schmitz-Rode T, Steinseifer U et al. 2014. Miniaturization: The clue to clinical application of the artificial placenta. *Artif Organs*. 38(3):208-214.

52. Rochow N, Manan A, Wu WI, Fusch G, Monkman S, Leung J, Chan E, Nagpal D, Predescu D, Brash J et al. 2014. An integrated array of microfluidic oxygenators as a neonatal lung assist device: In vitro characterization and in vivo demonstration. *Artif Organs*. 38(10):856-866.
53. Major TC, Handa H, Annich GM, Bartlett RH. 2014. Development and hemocompatibility testing of nitric oxide releasing polymers using a rabbit model of thrombogenicity. *Journal of biomaterials applications*. 29(4):479-501.
54. Reynolds MM, Annich GM. 2011. The artificial endothelium. *Organogenesis*. 7(1):42-49.
55. Walsh-Sukys MC, Bauer RE, Cornell DJ, Friedman HG, Stork EK, Hack M. 1994. Severe respiratory failure in neonates: Mortality and morbidity rates and neurodevelopmental outcomes. *The Journal of pediatrics*. 125(1):104-110.
56. Callaghan JC, Angeles J, Boracchia B, Fisk L, Hallgren R. 1963. Studies of the first successful delivery of an unborn lamb after 40 minutes in the artificial placenta. *Canadian journal of surgery Journal canadien de chirurgie*. 6:199-206.
57. Chamberlain G. 1968. An artificial placenta: The development of an extracorporeal system for maintenance of immature infants with respiratory problems. *American journal of obstetrics and gynecology*. 100(5):615-626.

58. Alexander DP, Britton HG, Nixon DA. 1968. Maintenance of sheep fetuses by an extracorporeal circuit for periods up to 24 hours. American journal of obstetrics and gynecology. 102(7):969-975.
59. Griffith BP, Borovetz HS, Hardesty RL, Hung TK, Bahnson HT. 1979. Arteriovenous ecmo for neonatal respiratory support. A study in perigestational lambs. The Journal of thoracic and cardiovascular surgery. 77(4):595-601.
60. Reoma JL, Rojas A, Kim AC, Khouri JS, Boothman E, Brown K, Grotberg J, Cook KE, Bartlett RH, Hirschl RB et al. 2009. Development of an artificial placenta i: Pumpless arterio-venous extracorporeal life support in a neonatal sheep model. Journal of pediatric surgery. 44(1):53-59.
61. Miura Y, Matsuda T, Funakubo A, Watanabe S, Kitanishi R, Saito M, Hanita T. 2012. Novel modification of an artificial placenta: Pumpless arteriovenous extracorporeal life support in a premature lamb model. Pediatric research. 72(5):490-494.

FIGURE 1




FIGURE LEGEND

Figure 1. Artificial placenta circuit configurations. (A) Pumpless arterio-venous extracorporeal lung support (AV-ECLS): blood inflow to the extracorporeal circuit is performed by cannulation of umbilical arteries; the oxygenated blood is then returned to EPT infant through the umbilical vein. (B) Pump-assisted veno-venous extracorporeal lung support (VV-ECLS): blood is drained from central vein(s) (e.g. internal jugular vein); the oxygenated blood is then returned to EPT infant through the umbilical vein.

TABLE 1 - Overview of early experiments performed with several AP models

Reference	Year	Model	Circuit	Oxygenator	Pump	Submersion	Survival
25	1958	Human	AV-ECLS	Rotating film	Yes	Yes	5-12 hours
26	1961	Lamb	VV-ECLS	Rotating disc	Yes	Yes	8-19 hours
27	1962	Dog	VV-ECLS	Rotating disc	Yes	Yes	2.5 hours
28	1962	Piglet	AV-ECLS	Rotating disc film	Yes	Yes	8 hours
56	1963	Lamb	AV-ECLS	Rotating disc film	Yes	Yes	40 minutes
29	1964	Lamb	AV-ECLS	Rotating disc film	Yes	No	1 hour
30	1964	Dog	AV-ECLS	Membrane	Yes	-	2-5 hours
12	1965	Lamb	AV-ECLS	Rotating disc film	No	Yes	0.3-3 hours
57	1968	Human	-	Coiled membrane	Yes	-	1.5-5 hours
58	1968	Lamb	AV-ECLS	Rotating disc film	Yes	Yes	24 hours
32	1969	Lamb	AV-ECLS	Silicone coiled	Yes	Yes	4-55 hours

TABLE 1 (continued)

Reference	Year	Model	Circuit	Oxygenator	Pump	Submersion	Survival
59	1979	Lamb	AV-ECLS	Microchannel membrane	Yes	No	-
38	1987	Goat	AV-ECLS	Silicone hollow fiber	Yes	Yes	Up to 165 hours
39	1989	Goat	AV-ECLS	Silicone hollow fiber	Yes	Yes	Up to 236 hours
40	1993	Goat	AV-ECLS	Silicone hollow fiber	Yes	Yes	Up to 542 hours
46	1998	Goat	AV-ECLS	Polyolefin hollow fiber	Yes	Yes	Up to 237 hours

TABLE 2 - Overview of recent experiments performed with several AP models

Reference	Year	Model	Circuit	Oxygenator	Pump	Submersion	Survival
60	2009	Lamb	AV-ECLS	Hollow fiber	No	No	4 hours
50	2011	Lamb	AV-ECLS	Polypropylene	No	No	3 hours
61	2012	Lamb	AV-ECLS	Membrane	No	Yes	Up to 30 hours
48	2012	Lamb	VV-ECLS	Polypropylene	Yes	Yes	24 hours
49	2013	Lamb	VV-ECLS	Polypropylene	Yes	No	70 hours
52	2014	Piglet	AV-ECLS	Microfluidic	No	No	4 hours
51	2014	Lamb	AV-ECLS	Polypropylene	No	No	6 hours
16	2015	Lamb	VV-ECLS	Polypropylene	Yes	No	Up to 1 week

ANEXOS

I - 2018 Update on “*Artificial Placenta: Recent Advances and Potential Clinical Applications*”

II - Editorial “*The Artificial Placenta: is Clinical Translation Next?*”

III - Citações do artigo “*Artificial Placenta: Recent Advances and Potential Clinical Applications*”

IV - Normas de publicação da revista científica *Pediatric Pulmonology*

V - Agradecimentos

2018 UPDATE ON “ARTIFICIAL PLACENTA: RECENT ADVANCES AND POTENTIAL CLINICAL APPLICATIONS”

In 2016, Miura and colleagues modified membranous oxygenators to test the hypothesis of a parallelized AP circuit¹, and compared it with the single-circuit AP. Parallelization of the AP system successfully decreased its resistance and prolonged low-weight lamb fetuses' survival up to 64 hours. Moreover, significantly low blood lactate levels were registered¹. Further studies under a parallelized circuit were conducted, with lamb fetuses being kept in a physiologically stable condition for periods of 48 hours² and 1 week³. However, white-matter injury was reported in two out of five animals, in each experiment^{1,2}.

Aiming the study of cerebral perfusion and oxygenation, researchers from Michigan University ECLS Laboratory successfully maintained fourteen lambs under venovenous AP support for up to 92 hours⁴. Despite the fact that this particular circuit configuration presents high risk for cerebral hypoperfusion, spectroscopy and carotid arterial flow suggested that brain oxygen delivery was preserved⁴ and that it poorly correlated with systemic oxygen saturation⁴. Further studies are necessary to assess white matter injury, and address the issue of intraventricular hemorrhage, even though necropsy evaluations showed no evidence of intracranial hemorrhage in their study⁴.

This group of researchers recently concluded that intratracheal perfluorocarbon instillation during AP support prevents lung injury and maximizes lung development, in comparison with the previously preferred approach with amniotic fluid and tracheal occlusion⁵. Increased surfactant production was also reported⁵. With further development, the VV-ECLS may offer a feasible alternative for EPT infants.

In 2017, Flake and colleagues presented a unique AP system that incorporates a pumpless oxygenator through an umbilical vascular interface kept within a closed fluid circuit, the polyethylene Biobag⁶. This model, which resembles the womb environment in shape and size, allowed eight fetal lambs to grow in a temperature-controlled, near-sterile environment, breathing an amniotic-like fluid⁶. Swallowing an electrolyte solution improved fetal fluid homeostasis and provided an additional route for nutrition. Furthermore, since there was continuous fluid exchange, this configuration solved the problem of gross fluid contamination and infection⁶.

In this study, double umbilical artery and single umbilical vein cannulation were preferred over carotid use, preserving a length of native umbilical cord between the cannula tips and the abdominal wall. This approach allowed vascular adaptation to pressure changes across the AV shunt⁷ and was proven to optimize circuit flow dynamics and stability, since it closely relates to the placental physiology in comparison to other cannulation methods⁷.

Remarkably, Flake et al. reported successful transition to air breathing after AP support in two lambs that were connected to the circuit for 21 and 29 days, respectively⁸. The animals presented satisfying responses to feeding and normal somatic growth⁸. One did not survive a pyelonephritis after 12 days of independent life, whereas the other reportedly surpassed three months survival with apparently normal neurological outcomes⁸.

These results are historical and superior to all previous attempts of EPT lambs' ECLS in both duration and physiologic well-being. Researchers will carry on evaluating and refining this AP system, since it needs to be downsized for human infants, who are one-third the size of the experimental fetus lambs. The first clinical trial is expected to occur within five years.

REFERENCES

1. Miura Y, Matsuda T, Usuda H, Watanabe S, Kitanishi R, Saito M, Hanita T, Kobayashi Y. 2016. A parallelized pumpless artificial placenta system significantly prolonged survival time in a preterm lamb model. *Artif Organs*. 40(5):E61-68.
2. Usuda H, Watanabe S, Miura Y, Saito M, Musk GC, Rittenschober-Bohm J, Ikeda H, Sato S, Hanita T, Matsuda T et al. 2017. Successful maintenance of key physiological parameters in preterm lambs treated with ex vivo uterine environment therapy for a period of 1 week. *Am J Obstet Gynecol*. 217(4):457.e451-457.e413.
3. Miura Y, Usuda H, Watanabe S, Woodward E, Saito M, Musk GC, Kallapur SG, Sato S, Kitanishi R, Matsuda T et al. 2017. Stable control of physiological parameters, but not infection, in preterm lambs maintained on ex vivo uterine environment therapy. *Artif Organs*. 41(10):959-968.
4. El-Sabbagh AM, Gray BW, Shaffer AW, Bryner BS, Church JT, McLeod JS, Zakem S, Perkins EM, Shellhaas RA, Barks JDE et al. 2017. Cerebral oxygenation of premature lambs supported by an artificial placenta. *Asaio j*.
5. Church JT, Perkins EM, Coughlin MA, McLeod JS, Boss K, Bentley JK, Hershenson MB, Rabah R, Bartlett RH, Mychaliska GB. 2018. Perfluorocarbons prevent lung injury and promote development during artificial placenta support in extremely premature lambs. *Neonatology*. 113(4):313-321.

6. Partridge EA, Davey MG, Hornick MA, McGovern PE, Mejaddam AY, Vrecenak JD, Mesas-Burgos C, Olive A, Caskey RC, Weiland TR et al. 2017. An extra-uterine system to physiologically support the extreme premature lamb. *Nature Communications*. 8:15112.
7. Hornick MA, Davey MG, Partridge EA, Mejaddam AY, McGovern PE, Olive AM, Hwang G, Kim J, Castillo O, Young K et al. 2018. Umbilical cannulation optimizes circuit flows in premature lambs supported by the extra-uterine environment for neonatal development (extend). *J Physiol*.
8. Partridge EA, Davey MG, Hornick MA, Flake AW. 2017. An extrauterine environment for neonatal development: Extending fetal physiology beyond the womb. *Semin Fetal Neonatal Med*. 22(6):404-409.

The Artificial Placenta: Is Clinical Translation Next?

George B. Mychaliska, MD*

Despite significant advances in the treatment of prematurity including antenatal steroids, advanced mechanical ventilation strategies, and exogenous surfactant, the mortality and morbidity remain high for these vulnerable infants. In particular, the mortality and morbidity of extremely low gestational age newborns (ELGANs) defined as <28 weeks estimated gestational age (EGA), is extremely high.¹ A radical paradigm shift in the treatment of extreme prematurity would be to recreate the intra-uterine environment using an extracorporeal artificial placenta (AP).

In this issue, Metelo-Coimbra and Roncon-Albuquerque² review recent advances in the field and assess barriers to clinical translation. From the outset, it should be acknowledged that although there is a large number of premature births worldwide (defined as <37 weeks EGA), outcomes have substantially improved for infants >28 weeks EGA. Apart from some specific congenital anomalies like congenital diaphragmatic hernia, the AP is intended for the treatment of ELGANS who experience the most complications of prematurity and whose outcome remains poor.

The authors focus on lung immaturity and provide substantial evidence of the iatrogenic effects of mechanical ventilation on both lung injury and the deleterious cardiovascular effects.² It is worth noting that although ELGANS are predisposed to interventricular hemorrhage (IVH) given their immature germinal matrix, mechanical ventilation has been implicated in the pathogenesis of IVH by increasing intrathoracic and intracranial pressure with every breath.³ The ELGANS who are never subjected to positive airway pressure have fewer complications of prematurity. An appreciation of the deleterious effects of mechanical ventilation and high oxygen concentrations on premature lungs has led to a dramatic shift toward less invasive ventilator strategies for premature infants. Although this strategy appears promising for some patients,⁴ there is still a subset of ELGANS that cannot maintain adequate gas exchange with the most invasive ventilator strategies.

Although the pulmonary system is critical to initial survival and long-term pulmonary morbidity is high among survivors, there are other significant complications of ELGANs that warrant consideration. Predictable and unsolved complications associated with prematurity include neurologic injury (IVH, white matter injury), necrotizing enterocolitis (NEC), retinopathy of prematurity (ROP), and sepsis. Our current inability to prevent these complications relates both to organ immaturity and conventional treatment strategies such as positive pressure ventilation, that have historically been developed for full term infants. To potentially solve these problems, the AP must not only recreate the fetal milieu and provide life-sustaining functions such as adequate gas exchange and fetal hemodynamic stability, but it should also protect against organ trauma and allow the normal developmental pathways to occur.

An AP may appear far beyond the reach of modern science, but the idea of creating a life support system to maintain growing fetuses in a womb-like environment with extracorporeal support was first investigated 60 years

Section of Pediatric Surgery, Department of Surgery, Fetal Diagnosis and Treatment Center, University of Michigan Medical School, C.S. Mott Children's Hospital, 1540 E. Medical Center Drive, SPC 4211, Ann Arbor, Michigan 48109.

Conflict of Interest: None.

Funding source: NIH, Number 1R01HD073475-01A1.

*Correspondence to: George B. Mychaliska, MD, Section of Pediatric Surgery, Department of Surgery, Fetal Diagnosis and Treatment Center, University of Michigan Medical School, C.S. Mott Children's Hospital, 1540 E. Medical Center Drive, SPC 4211, Ann Arbor, MI 48109. E-mail: mychalisk@med.umich.edu

Received 20 February 2016; Accepted 6 March 2016.

DOI 10.1002/ppul.23412

Published online 19 April 2016 in Wiley Online Library (wileyonlinelibrary.com).

ago! Metelo-Coimbra and Roncon-Albuquerque provide a succinct review of AP terminology and history of milestones.² Since fetuses normally develop with extracorporeal support, it should perhaps not surprise us that researchers were drawn to this concept shortly after the successful introduction of cardio-pulmonary bypass. For historians of science, it is noteworthy that researchers were on the right path, but got derailed many times due to the state of biomedical technology and insufficient knowledge of the physiology of premature infants. In my view, the history of the development of the AP is marked by many experimental failures with episodic successes. Despite incremental success, many research groups abandoned this work as progress was being made with antenatal steroids, exogenous surfactant, advanced mechanical ventilation strategies, and ECMO for term and near-term infants.

An appreciation of the unsolved problems of extreme prematurity coupled with recent advances detailed by the authors² has led to a resurgence of work on the AP. The fetal lamb is the best model, and the lamb gestational age which corresponds to ELGAN lungs is 118 days gestation (term = 145 days). Although the AP is promising and has the potential to radically change the treatment of prematurity, several obstacles remain. As the authors point out,² the first issue is the most effective ECLS configuration. A simple pumpless AV-ECLS circuit utilizing the umbilical vessels is appealing, but our experience demonstrated only short-term survival and declining cardiac function.⁵ Despite cannulation of the umbilical arteries to the sheep aorta (to obviate vessel spasm) and adding a pump, matching extracorporeal flow to systemic flow is very difficult (the native placenta does this automatically). In addition, given the tortuosity, size, and spasm associated with human umbilical arteries, we transitioned to a pump-driven VV-ECLS model with inflow via the umbilical vein and outflow via the jugular vein. This approach provides 7 days of support with excellent gas exchange and hemodynamic stability.⁶ With current technology, we believe this strategy is clinically translatable to extremely premature infants.

As mentioned previously, the AP strategy will require long-term support (2–4 weeks in humans) and demonstration that organs are maturing and protected from trauma. This corresponds to 10–14 days in the 118 day lamb model. As such, an in-depth study of lung development, long-term support, and weaning to a ventilator and air breathing will be required. A crucial aspect of lung development will be the airway strategy during AP support. In our early work, the fetal lambs were submerged in a warmed “amniotic bath” effectively recreating the intrauterine environment.^{5,7} While appealing in some regards, there are infection and patient access issues with this approach. More recently, we have been intubating the fetal lambs, filling them with amniotic fluid or Perflubron and either capping the endotracheal tube or

maintaining 5–8 cm H₂O pressure. It is possible to harness the power of mechanotransduction with this approach and possibly accelerate lung growth.^{8,9}

Apart from lung development which is crucial during AP support, other vulnerable premature organs need to be examined. Brain perfusion, function, and development are critical to understand. Although the sheep is not a good model for IVH, brain physiology and evidence of white matter injury can be assessed. With a high incidence of NEC in premature infants, optimal nutrition and perfusion of the gastrointestinal system warrants investigation. Long term survivors of the AP should be examined for evidence of retinopathy of prematurity. Lastly, renal and hepatic function should be addressed.

As a general rule, extracorporeal support is reserved for infants ≥ 34 weeks EGA due to a higher rate of IVH in extremely premature infants. The authors point out the feasibility of “preemie ECMO” in infants from 29 to 33 weeks,² but ELGANs would have prohibitively high rates of IVH. As such, a critical barrier to clinical application will be the development of non-thrombogenic surfaces that will obviate the need for anti-coagulation.^{10,11} Lastly, clinical application will require a clinical prognostication tool to select premature infants at the highest risk for mortality on the first day of life.¹²

Given recent advances and ongoing work, we believe that the AP will be used in ELGANs in the next 5 years.

REFERENCES

1. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, Hale EC, Newman NS, Schibler K, Carlo WA, et al. Neonatal outcomes of extremely preterm infants from the NICHD Neonatal Research Network. *Pediatrics* 2010;126:443–456.
2. Metelo-Coimbra C, Roncon-Albuquerque R, Jr. Artificial placenta: recent advances and potential clinical applications. *Pediatr Pulmonol*. doi: 10.1002/ppul.23401.
3. Aly H, Hammad TA, Essers J, Wung JT. Is mechanical ventilation associated with intraventricular hemorrhage in preterm infants? *Brain Dev* 2012;34:201–205.
4. Schmolzer GM, Kumar M, Pichler G, Aziz K, O'Reilly M, Cheung PY. Non-invasive versus invasive respiratory support in preterm infants at birth: systematic review and meta-analysis. *BMJ* 2013;347:f5980.
5. Reoma J, Rojas A, Kim A, Khouri J, Boothman E, Brown K, Grotberg J, Cook K, Bartlett R, Hirschl R, et al. Development of an artificial placenta I: pumpless arterio-venous extracorporeal life support in a neonatal sheep model. *J Pediatr Surg* 2009;44:53–59.
6. Bryner B, Gray B, Perkins E, Davis R, Hoffman H, Barks J, Owens G, Bocks M, Rojas-Pena A, Hirschl R, et al. An extracorporeal artificial placenta supports extremely premature lambs for 1 week. *J Pediatr Surg* 2015;50:44–49.
7. Gray BW, El-Sabbagh A, Rojas-Pena A, Kim AC, Gadepali S, Koch KL, Capizzani TR, Bartlett RH, Mychaliska GB. Development of an artificial placenta IV: 24 hour venovenous extracorporeal life support in premature lambs. *ASAIO J* 2012;58:148–154.
8. Mychaliska G, Bryner B, Dechert R, Kreutzman J, Becker M, Hirschl R. Safety and efficacy of perflubron-induced lung growth in neonates with congenital diaphragmatic hernia: results of a prospective randomized trial. *J Pediatr Surg* 2015;50:1083–1087.

9. Shue EH, Miniati D, Lee H. Advances in prenatal diagnosis and treatment of congenital diaphragmatic hernia. *Clin Perinatol* 2012;39:289–300.
10. Major TC, Brant DO, Burney CP, Amoako KA, Annich GM, Meyerhoff ME, Handa H, Bartlett RH. The hemocompatibility of a nitric oxide generating polymer that catalyzes S-nitrosothiol decomposition in an extracorporeal circulation model. *Biomaterials* 2011;32:5957–5969.
11. Major TC, Brant DO, Reynolds MM, Bartlett RH, Meyerhoff ME, Handa H, Annich GM. The attenuation of platelet and monocyte activation in a rabbit model of extracorporeal circulation by a nitric oxide releasing polymer. *Biomaterials* 2010;31:2736–2745.
12. Reid S, Bajuk B, Lui K, Sullivan EA. Comparing CRIB-II and SNAPPE-II as mortality predictors for very preterm infants. *J Paediatr Child Health* 2015;51:524–528.

CITAÇÕES DO ARTIGO “ARTIFICIAL PLACENTA: RECENT ADVANCES AND POTENTIAL CLINICAL APPLICATIONS”

1. Mychaliska G. 2016. The artificial placenta: Is clinical translation next? *Pediatric pulmonology*. 51(6):557-559.
2. Bird SD. 2017. Artificial placenta: Analysis of recent progress. *Eur J Obstet Gynecol Reprod Biol*. 208:61-70.
3. Parga JJ, Garg M. 2017. Extracorporeal membrane oxygenation in neonates: History and future directions. *NeoReviews*. 18(3):e166.
4. te Pas AB. 2017. Improving neonatal care with technology. *Frontiers in Pediatrics*. 5:110.
5. Mazumdar Bolanos M. 2017. Electrolysis-based system for generation and delivery of oxygen to microfluidic oxygenator unit for preterm neonates with respiratory distress syndrome. [MacSphere - McMaster University Libraries Institutional Repository]: McMaster University.
6. Surate Solaligue DE, Rodriguez-Castillo JA, Ahlbrecht K, Morty RE. 2017. Recent advances in our understanding of the mechanisms of late lung development and bronchopulmonary dysplasia. *Am J Physiol Lung Cell Mol Physiol*. 313(6):L1101-L1153.

Pediatric Pulmonology

© Wiley Periodicals, Inc.



Edited By: Thomas Murphy

Impact Factor: 2.758

ISI Journal Citation Reports © Ranking: 2016: 19/121 (Pediatrics); 26/59 (Respiratory System)

Online ISSN: 1099-0496

Author Guidelines

- SCOPE OF JOURNAL
- PERMISSIONS
- AUTHOR RESOURCES
- ENGLISH LANGUAGE SERVICES
- ELECTRONIC SUBMISSION OF MANUSCRIPTS
- MANUSCRIPT GUIDELINES
 - Original Research Articles
 - Reviews/State of the Art Papers
 - Case Reports
 - Editorials (Commentaries)
 - Letters to the Editor
- PRIOR TO SUBMITTING
- COMPONENTS OF ARTICLES/FILE PREPARATION
 - Main Document
 - Title Page
Summary/Abstract
Acknowledgements
Informed Consent
References

Keywords
Abbreviations
Drug Names
Eponyms
Formatting Specific to Original Research Articles

- [Tables](#)
- [Images](#)
- [Online Supporting Information](#)

- [POLICIES/DISCLOSURE STATEMENTS](#)
 - [Conflict of Interest](#)
 - [Experimental and Publication Ethics](#)
 - [Plagiarism](#)
 - [Prior Publication](#)
 - [Clinical Trials](#)

- [PEER REVIEW PROCESS](#)
- [FAST TRACK REVIEW](#)
- [SUBMISSIONS FROM EDITORS AND EDITORIAL BOARD MEMBERS](#)
- [AUTHOR CHARGES](#)
- [MANUSCRIPTS ACCEPTED FOR PUBLICATION](#)
 - [Online Open](#)
 - [Copyright Transfer Agreement](#)

- [PROOFS](#)
- [REPRINTS](#)
- [APPEALS PROCESS](#)
- [PRODUCTION QUESTIONS](#)
- [QUESTIONS ABOUT YOUR SUBMISSION](#)
- [CONTACT THE EDITOR-IN-CHIEF](#)

SCOPE OF JOURNAL

Pediatric Pulmonology publishes the results of original clinical or laboratory research, state of the art reviews, exceptionally instructive or unique case reports, and letters to the Editor (and responses), pertaining to the specialty.

Reports on meetings, conferences and symposia may be published after consultation with the Publisher and the Editor-in-Chief.

Preliminary brief communications will be considered if the articles contain information which would be considered a major breakthrough in the field.

We do not publish research funded by tobacco companies.

As the field is continually evolving, our Journal has seen an increase in the number of submissions over the past few years, and, as a result, our rejection rate is climbing.

PERMISSIONS

No material published in Pediatric Pulmonology may be reproduced or published elsewhere without the written permission of the publisher and the author. To request permission to reproduce an article, in part, or in whole, click here to for the [Permissions Page \(http://onlinelibrary.wiley.com/journal/10.1002/\(ISSN\)1099-0496/homepage/Permissions.html\)](http://onlinelibrary.wiley.com/journal/10.1002/(ISSN)1099-0496/homepage/Permissions.html)

AUTHOR RESOURCES

For additional tools visit [Author Services \(http://authorservices.wiley.com/bauthor/default.asp\)](http://authorservices.wiley.com/bauthor/default.asp) - an enhanced suite of online tools for WileyOnlineLibrary journal authors, featuring Article Tracking, E-mail Publication Alerts and Customized Research Tools.

ENGLISH LANGUAGE SERVICES

The Editors reserve the right to return any manuscript that is not in acceptable English. Translations from another language will not be provided by the Editorial Office. Authors from countries in which English is not the primary language should have their manuscript reviewed and corrected by an English language service before submission. To read more about our policy, and to view a list of editing services, visit: http://authorservices.wiley.com/bauthor/english_language.asp (http://authorservices.wiley.com/bauthor/english_language.asp).

GUIDELINES FOR COVER SUBMISSIONS

If you would like to send suggestions for artwork related to your manuscript to be considered to appear on the cover of the journal, please [follow these general guidelines. \(http://olabout.wiley.com/WileyCDA/Section/id-828302.html\)](http://olabout.wiley.com/WileyCDA/Section/id-828302.html)

ELECTRONIC SUBMISSION OF MANUSCRIPTS

If you are familiar with our guidelines, click [here \(http://mc.manuscriptcentral.com/ppul\)](http://mc.manuscriptcentral.com/ppul) to login to your ScholarOne account to submit your manuscript. If you do not have an account, click on “Register Here” to establish one.

MANUSCRIPT GUIDELINES

We accept submissions of the following types of articles. Please note the specific guidelines for each type:

Original Research Articles

Original Research Articles should follow the standard structure of abstract, introduction, methods, results, discussion, and references, and may include up to six tables and/or images when appropriate. Original Research Articles should be limited to 3,500 words (not including the abstract or references). The abstract should not exceed 250 words, and references should be limited to forty (40).

Reviews/State of the Art Papers

Editors generally commission Reviews and State of the Art papers, but uninvited submissions are also welcome, particularly if the submission outlines an important and topical subject with a focus on recent advances. Reviews should be limited to 4,000

words, while State of the Art papers should be limited to 5,000 words (not including the abstract or references). We ask that the abstracts for these manuscript types do not exceed 250 words. There is no set limit on images, tables, or references for these types of manuscripts.

Case Reports

Pediatric Pulmonology will review case report manuscripts that present unique, paradigm-changing, or novel accounts of infantile or childhood disorders. Priority for selection for publication will be given to the following categories:.

1. Novel therapies and outcomes for cystic fibrosis
2. Novel disorders or outcomes of ChILD, NEHI, ABCA3 disorders, and surfactant disorders
3. Novel congenital malformations
4. Novel genetic disorders
5. Novel therapies or outcomes for other disorders

There is not a related format for a case series. Manuscripts of this nature will be treated as original articles or reviews and will compete with other manuscripts in these categories.

Case reports should be concise (a maximum of 1,000 words, not including the abstract or references), and contain a maximum of two images and/or tables. The summary/abstract should not exceed 100 words. Case Reports should contain no more than five (5) references. Authorship of case reports shall be limited to three (3). Physicians who participated in the care, but did not contribute to the writing of the manuscript may be listed under acknowledgements. Informed consent must be documented. Authors should note that most accepted Case Reports will be published online only, and not in a print edition.

Editorials (Commentaries)

Editors and members of the Editorial Board may make editorial comments on individual articles or on a group of articles published in the same issue. Editorials (including pro/con debates) from authors who are not part of the editorial team are also welcome as submissions to the Journal. These narrowly focused articles should discuss an article that was recently published, or that is soon to be published. The commentary should discuss specific issues within a subject area rather than the whole field, while explaining the implications of the article and putting it in context. Opinions must be factually based. These types of manuscripts should be limited to 1,500 words (not including the abstract or references). There is no limitation on the number of tables, images or references for these types of manuscripts.

Letters to the Editors

We encourage letters that offer criticism of published material in an objective, constructive, and educational manner conducive to further exchanges. Such letters will only be considered if they are in reference to an article published within the previous six months. Letters may also discuss matters of general interest pertaining to the field of pediatric pulmonology, or may consist of brief reports of truly unique cases. Note that we do not publish original, previously unpublished data as letters. If appropriate, a copy will be sent to the author(s) referred to in the letter, so that they may respond. Letters to the Editor should not exceed 1,000 words (not including the abstract or references), and may contain a small table or single image. Letters should contain no more than five (5) references.

PRIOR TO SUBMITTING

Prior to submitting a manuscript through ScholarOne (<http://mc.manuscriptcentral.com/ppul>), prepare the text and images according to the instructions found below. You may enter and exit the manuscript submission process at the completion of each step, and you may save an unfinished draft in the system to work on later. However, once you submit your manuscript through the system, you will not be able to access it for editing. If you have any questions about this process please contact us at edsupport@wiley.com (<mailto:edsupport@wiley.com>).

We recommend all authors familiarize themselves with the International Committee of Medical Journal Editors: Uniform Requirements for Manuscripts submitted to Biomedical Journals. Ann Intern Med 1997;126:36-47. The complete text of the document can be found online at www.icmje.org (<http://www.icmje.org>).

COMPONENTS OF ARTICLES/FILE PREPARATION

Please make note of the following when preparing your submission:

Main Document

All manuscript types must include a title page, abstract, text and references in the Main Document. Standard, double-spaced manuscript format, in 12 point font is requested. Number all pages consecutively.

Title page: The title should be brief (no more than 100 characters in length including spaces) and useful for indexing. All authors' names with highest academic degree, affiliation of each, but no position or rank, should be listed. For cooperative studies, the institution where research was primarily done should be indicated. In a separate paragraph, specify grants, other financial support received, and the granting institutions (grant number(s) and contact name(s) should be indicated on the title page). If support from manufacturers of products used is listed, assurances about the absence of bias by the sponsor and principal author must be given. Identify meetings, if any, at which the paper was presented. The name, complete mailing address, telephone number, fax number, and e-mail address of the person to whom correspondence and reprint requests are to be sent must be included. Keywords should also be noted on the title page. For usage as a running head, provide an abbreviated title (maximum 50 characters) on the bottom of the title page.

Summary/Abstract: In accordance with the structure of the article, with or without separate headings, outline the objectives, working hypothesis, study design, patient-subject selection, methodology, results (including numerical findings) and conclusions. The Summary should not exceed the word counts outlined above. If abbreviations are used several times, spell out the words followed by the abbreviations in parentheses.

Acknowledgements: Technical assistance, advice, referral of patients, etc. may be briefly acknowledged at the end of the text under "Acknowledgements."

Informed Consent: Informed consent statements, if applicable, should be included in the Methods section.

References/citations: References may be included at the end of your text, or uploaded as a separate file. Ensure your references are up to date, and include a critical selection from the world literature. References should be prepared according to CSE (Council of

Science Editors) citation-sequence style. Refer to the *Scientific Style and Format: The CSE Manual for Authors, Editors, and Publishers*, 8th edition (University of Chicago Press). Start the listing on a new page, double-spaced throughout.

Number the references in the sequence in which they first appear in the text, listing each only once even though it may be cited repeatedly.

When citing a reference in the text, the style advocated by CSE suggests numbers appear in superscript, and appear before punctuation marks (commas or periods). In the **citation-sequence** system, sources are numbered by order of reference so that the first reference cited in the paper is ¹, the second ², and so on. If the numbers are not in a continuous sequence, use commas (with no spaces) between numbers. If you have more than two numbers in a continuous sequence, use the first and last number of the sequence joined by a hyphen, for example ^{2,4,6-10}.

In the references, list the first ten authors of the cited paper. If there are more than ten authors, list the first 10 authors followed by 'et al'.

Journals' names should be shown by their abbreviated title in *Index Medicus*.

Manuscripts in preparation or submitted for publication are not acceptable references. If a manuscript "in press" is used as a reference, a copy of it must be provided with your submission.

Sample references:

Standard journal article

Landau IL, Morgan W, McCoy KS, Taussig LM. Gender related differences in airway tone in children. *Pediatr Pulmonol* 1993;16:31-35.

Book with authors

Voet D, Voet JG. 1990. *Biochemistry*. New York: John Wiley & Sons. 1223 p.

Book with editors

Coutinho A, Kazatch Kine MD, editors. *Autoimmunity physiology and disease*. New York. Wiley-Liss; 1994. 459 p.

Chapter from a book

Hausdorf G. Late effects of anthracycline therapy in childhood: evaluation and current therapy. In: Bricker JT, Green DM, D'Angio GJ, editors. *Cardiac toxicology after treatment for childhood cancer*. New York: Wiley-Liss; 1993. p 73-86.

For a book reference only include the page numbers that have direct bearing on the work described.

Keywords: On the title page, supply a minimum of 3 to 5 keywords, exclusive of words in the title of the manuscript. A guide to medical subject heading terms used by PubMed is available at <http://www.nlm.nih.gov/mesh/MBrowser.html> (<http://www.nlm.nih.gov/mesh/MBrowser.html>).

Abbreviations: Define abbreviations when they first occur in the manuscript and from there on use only the abbreviation. Whenever standardized abbreviations are available use those. Use standard symbols with subscripts and superscripts in their proper place.

Drug names: Use generic names. If identification of a brand name is required, insert it in parentheses together with the manufacturer's name and address after the first mention of the generic name.

Eponyms: Eponyms (diseases or biologic entities named for persons) should not be used when standard descriptive terminology is available. Examples include club cells (formerly known as Clara cells); and granulomatosis with polyangiitis (formerly known as Wegener's granulomatosis). It is permissible to use the eponym in parenthesis at the first mention of the term in cases in which the eponym is still in common use.

Formatting Specific to Original Research Articles: Divide article into: Title Page, Summary/Abstract, Introduction, Materials and Methods, Results, Discussion, and References, starting each section on a new page. All methodology and description of experimental subjects should be under Materials and Methods; results should not be included in the Introduction. Please ensure the following appears in the appropriate section of your manuscript:

- a concise introductory statement outlining the specific aims of the study and providing a discussion of how each aim was fulfilled;
- a succinct description of the working hypothesis;
- a detailed explanation of assumptions and choices made regarding study design and methodology;
- a description of the reasons for choosing the type and number of experimental subjects (patients, animals, controls) and individual measurements; if applicable, information about how and why the numbers may differ from an ideal design (e.g., the number required for achieving 90% confidence in eliminating Type II error);
- specifics about statistical principles, techniques and calculations employed and, if applicable, methods for rejecting the null hypothesis;
- a concise comparison of the results with those of conflicting or confirmatory studies in the literature;
- a brief summary of the limitations of the scientific methods and results; and
- a brief discussion of the implications of the findings for the field and for future studies.

- **Tables**

Tables should not be included in the Main Document, but submitted as a separate DOC or RTF file. Number tables with Arabic numbers consecutively and in order of appearance. Type each table double-spaced on a separate page, captions typed above the tabular material. Symbols for units should be used only in column headings. Do not use internal horizontal or vertical lines; place horizontal lines between table caption and column heading, under column headings, and at the bottom of the table (above the footnotes if any). Use footnote letters (a, b, c, etc.) in consistent order in each table. All tables should be referred to in the text. Do not submit tables as photographs and do not separate legends from tables.

- **Images**

Image files must be submitted in TIF or EPS (with preview) formats. Do not embed images in the Main Document. Number images with Arabic numbers and refer to each image in the text. The preferred form is 5 X 7 inches (12.5 X 17.5 cm). Print reproduction requires files for full color images to be in a CMYK color space.

Please note authors are encouraged to supply color images regardless of whether or not they are amenable to paying the color reproduction fees. Color images will be published online, while greyscale versions will appear in print at no charge to the author. See Author Charges below.

Journal quality reproduction requires grey scale and color files at resolutions yielding approximately 300 ppi. Bitmapped line art should be submitted at resolutions yielding 600-1200 ppi. These resolutions refer to the output size of the file; if you anticipate that your images will be enlarged or reduced, resolutions should be adjusted accordingly.

Lettering on images should be of a size and weight appropriate to the content and the clarity of printing must allow for legibility after reduction to final size. Labeling and arrows on images must be done professionally. Spelling, abbreviations, and symbols should precisely correspond to those used in the text. Indicate the stain and magnification of each photomicrograph. Photographs of recognizable subjects must be accompanied by signed consent of the subject of publication. Images previously published must be accompanied by the author's and publisher's permission.

Image legends should be brief, and included as a separate DOC file under the heading: "Image Legends." When borrowed material is used, the source of the image should be shown in parentheses after its legend, either by a reference number or in full if not listed under References.

Online Supporting Information

Additional non-essential material such as text, appendices, tables, images, video, and soundtrack files may be submitted for posting as supporting information to an article. The scientific value of such material should be evident. The material should be submitted simultaneously with the manuscript so that it may undergo peer review. In naming these files, please note the file names should be preceded by the letter "E." For example "E-table 1," "E-image 1," "E-text," etc.

Note that supporting online material is not typeset, nor proofread following the review process, so please ensure the material is accurate and free of typographical errors. Supporting material should be prepared in the same manner as the print material.

While supporting information does not appear in the print version, a notation is made that supporting material is available online.

Top of Page

POLICIES/DISCLOSURE STATEMENTS

We recognize the importance of developing the highest ethical standards and we are committed to ethical publication practice. For more information on the publisher's policies, please see Wiley-Blackwell Guidelines on Publication Ethics and Best Practices www.wiley.com/bw/publicationethics (http://exchanges.wiley.com/publishing-ethics_252.html). Of particular importance is the section on Research Misconduct, which includes data fabrication, falsification, plagiarism, and inappropriate image manipulation.

Authors who submit to Pediatric Pulmonology should take heed of the following:

Conflict of Interest: Authors must indicate at the time of submission any potential conflict of interest (particularly of a fiscal nature) that may have a perceived influence on the results of the research. The existence of such does not automatically preclude publication. A conflict of interest statement should appear in the Acknowledgment section. For further information on Conflict of Interest please visit www.icjme.org (<http://www.icmje.org/conflicts-of-interest/>).

Experimental and Publication Ethics: Studies involving human subjects must conform to the guiding principles of the World Medical Association Declaration of Helsinki (<http://www.wma.net/en/30publications/10policies/b3/>). Human subjects must have given informed consent and the study must have been approved by the Committee on Human Research at the author(s) institution(s) and a statement to this effect must appear in the Methods section of the submitted article. It is also important to document in the Methods section that consent has been obtained from older children and adolescents. Similarly, animal studies must be approved by an Institutional Animal Research Review Board and a statement to this effect must appear in the Methods section. In addition, details of anesthesia and euthanasia must appear in the Methods section.

Plagiarism: It should be noted that Pediatric Pulmonology employs a plagiarism detection system. By submitting your manuscript to this journal you accept that your manuscript may be screened for plagiarism against previously published works. CrossCheck is a multi-publisher initiative to screen published and submitted content for originality. Pediatric Pulmonology uses iThenticate software to detect instances of overlapping and similar text in submitted manuscripts. To find out more about CrossCheck visit <http://www.crossref.org/crosscheck.html> (<http://www.crossref.org/crosscheck.html>).

Prior Publication: Manuscripts submitted to Pediatric Pulmonology may not have been published in any part or form in another publication of any type, professional or lay, including electronic publications, the exception being abstracts of no more than 400 words. Any material available via PubMed or other electronic sources is considered to have been published. When a question arises, the Editor-in-Chief will determine what constitutes duplicate publication. If duplicate publication is confirmed, the Editor-in-Chief will initiate a discussion with the sponsoring institution and the authors.

It is the responsibility of submitting authors to inform the Editor-in-Chief of potentially overlapping or related data either in submitted manuscripts or papers in press, and such manuscripts should be appended to the submission. If there is significant overlap in data with previously published articles this should be addressed by the author in the "Authors Comments" section during the submission process. In particular, giving reasons why the new submission should be published. The editors reserve the right to determine whether or not publication is warranted.

For further information on redundant or duplicate publication, please visit <http://www.icmje.org> (<http://www.icmje.org/recommendations/browse/publishing-and-editorial-issues/overlapping-publications.html>).

Clinical Trials: We endorse the Consolidated Statement of Reporting Trials (CONSORT Statement) Lancet 2002;357:1191-1194 which may be accessed at www.consort-statement.org (<http://www.consort-statement.org/>). In accordance with ICMJE standards, all clinical trials must be registered with a database that is publicly

accessible such as <http://clinicaltrials.gov/> However, other free of charge public registries are acceptable. For further information, please visit <http://www.icmje.org/about-icmje/faqs/clinical-trials-registration/> (<http://www.icmje.org/about-icmje/faqs/clinical-trials-registration/>).

PEER REVIEW PROCESS

Upon submission, authors are encouraged to submit names of experts who they deem appropriate to review their paper. Authors may also indicate persons to whom they do not wish the manuscript sent for review. In most cases, articles will be reviewed by at least two authorities as well as the editorial staff to determine validity, significance, novelty, and potential impact on the field of contents and conclusions. The reviewers will be selected by the Editor-in-Chief, Associate Editors, and/or Editorial Board members. The selection will be made on the basis of expertise, impartiality, and equal distribution among the available experts, regardless of geographic origin of the manuscript or locations of the reviewers. Authors will be advised within the shortest possible time whether their paper is accepted, requires major or minor revisions, or is rejected. All necessary efforts will be made to ensure a timely review process.

Authors should note the time from submission to final decision can be shortened by a timely return of a revised manuscript when revision has been requested.

The Editor-in-Chief reserves the right to reject any submission deemed not suitable for the journal after an in-house review.

FAST TRACK REVIEW

If circumstances so warrant, a fast-track review of a paper may be requested in the cover letter. At the Editor-in-Chief's discretion, a fast-track review will be undertaken to expedite manuscripts that deserve rapid review and publication. Expedited peer review and publication is rare and is reserved for timely presentation of significant data. If fast-track review is requested in the cover letter, the corresponding author will be informed if expedited review has been granted or not.

SUBMISSIONS FROM EDITORS AND EDITORIAL BOARD MEMBERS

Pediatric Pulmonology strives to ensure that any submission from the Editor-in-Chief, Deputy Editor, the Associate Editors, or from a member of the journal's Editorial Board receives an objective and unbiased evaluation. This is achieved by assigning any research article submitted by the Editor-in-Chief, Deputy or Associate Editors to an impartial referee who can maintain the integrity of the review process. When appropriate, Pediatric Pulmonology may also utilize the services of Guest Editors who are familiar with the peer review processes and policies of the journal. Articles submitted by Editorial Board members undergo a blinded peer review process that is as stringent as for those authors who are not on the Board. All submitting authors are automatically blinded to all aspects of the review process.

AUTHOR CHARGES (AuthorCharges)

Should authors wish for manuscript images appear in color in the print edition, color reproduction fees will be charged to the authors. Current color reproduction fees are \$800 per figure. Authors do, however, have the option of submitting color images for online publication, and greyscale images for the print edition at no charge, and we encourage authors to do so. For information on color charges, please contact Production Editor, at [ppulprod @wiley.com](mailto:ppulprod@wiley.com)

MANUSCRIPTS ACCEPTED FOR PUBLICATION

The author identified as the formal corresponding author for the paper will receive an email prompting them to login into Author Services where via the Wiley Author Licensing Service (WALS) they will be able to complete a license agreement on behalf of all authors on the paper:

Online Open

OnlineOpen (<http://olabout.wiley.com/WileyCDA/Section/id-406241.html>) fulfills RCUK, Wellcome Trust, NIH, and other funder mandates. Authors can use OnlineOpen to make their article open access and freely available to all on Wiley Online Library. Wiley also immediately deposit OnlineOpen articles in PubMed Central and PMC mirror sites. In addition, authors of OnlineOpen articles are permitted to post the final, published PDF of their article on a website, institutional repository or other free public server, immediately on publication.

With OnlineOpen, the author, the author's funding agency, or the author's institution pays a fee to ensure that the article is made open access.

If the OnlineOpen option is selected the corresponding author will have a choice of the following Creative Commons License Open Access Agreements (OAA):

(<http://creativecommons.org/licenses/by-nc-nd/2.0/uk/>)Creative Commons Attribution License OAA (<http://creativecommons.org/licenses/by-nc/3.0/>)
(<http://creativecommons.org/licenses/by-nc-nd/2.0/uk/>)

- Creative Commons Attribution Non-Commercial -NoDerivs License OAA
(<http://creativecommons.org/licenses/by-nc-nd/2.0/uk/>)

If you select the OnlineOpen option and your research is funded by The Wellcome Trust and members of the Research Councils UK (RCUK) you will be given the opportunity to publish your article under a CC-BY license supporting you in complying with Wellcome Trust and Research Councils UK requirements. For more information on this policy and the Journal's compliant self-archiving policy please visit: <http://www.wiley.com/go/funderstatement>
(<http://www.wiley.com/go/funderstatement>)

For RCUK and Wellcome Trust authors click on the link below to preview the terms and conditions of this license:

Creative Commons Attribution License OAA (<http://creativecommons.org/licenses/by-nc/3.0/>)

Note to NIH Grantees: Pursuant to the NIH mandate, Wiley Blackwell will post the accepted version of contributions authored by NIH grant-holders to PubMed Central upon acceptance. This accepted version will be made publicly available 12 months after publication. For further information, see Wiley Blackwell's NIH Policy Statement.
(<http://www.wiley.com/WileyCDA/Section/id-321171.html>)

Copyright Transfer Agreement

If the OnlineOpen option is not selected, the corresponding author will be presented with the copyright transfer agreement (CTA) to sign on behalf of all authors.

To preview the terms and conditions of the open access agreements, or the copyright transfer agreement, please visit the FAQs hosted on Wiley Author Services. See http://authorservices.wiley.com/bauthor/faqs_copyright.asp (http://authorservices.wiley.com/bauthor/faqs_copyright.asp) and <http://www.wileyopenaccess.com/details/content/12f25db4c87/Copyright--License.html> (<http://www.wiley.com/WileyCDA/Section/id-321171.html>).

PROOFS

Following acceptance, the corresponding author will be alerted by e-mail to access galley proofs in web-based proofing system. Corrections should be returned within 48 hours of receipt, as delays in returning galley proofs cause delays in publication. Alterations should be kept to a minimum. Costs of extensive alterations to the galley proof will be billed to the authors. All statements in (or omissions from) published manuscripts are the responsibility of the authors who are asked to carefully review proofs prior to publication.

REPRINTS

Reprints may be ordered at: <https://caesar.sheridan.com/reprints/eorder/order> (<https://caesar.sheridan.com/reprints/eorder/order.php?DBS=A&PUID=10089&IUID=4440&UNDO=1&SERVICE=2&ACCTUID=75>).

APPEAL REQUESTS

Authors who wish to request reconsideration of a rejected manuscript should direct their query to the Editor in Chief, Tom Murphy, murphyeditorppul@hotmail.com (<mailto:murphyeditorppul@hotmail.com>) or the PPUL editorial office, ppuledoffice@wiley.com (ppuledoffice@wiley.com). Requests must include the manuscript ID and a detailed description of why the authors believe the paper should be reconsidered.

Appeal requests will be evaluated by the Editor-in-Chief to determine if an appeal will be permitted. These appeals will be approved on a very limited basis. If the appeal is allowed, instructions will be provided on how to resubmit your paper. Authors should not resubmit their rejected paper without prior approval from the Editors and the Editorial Office. Requests that are sent elsewhere will not be considered.

PRODUCTION QUESTIONS?

Contact Production Editor at ppulprod@wiley.com (<mailto:ppulprod@wiley.com>).

QUESTIONS ABOUT YOUR SUBMISSION?

Contact the PPUL editorial office at ppuledoffice@wiley.com (ppuledoffice@wiley.com).

CONTACT THE EDITOR-IN-CHIEF

Thomas Murphy, murphyeditorppul@hotmail.com (<mailto:murphyeditorppul@hotmail.com>).

[Top of Page](#)

AGRADECIMENTOS

O trabalho refletido nesta monografia não teria sido possível sem a colaboração, estímulo e empenho de diversas pessoas que, direta ou indiretamente, contribuíram para a sua realização, e às quais não poderia deixar de agradecer.

Ao Professor Doutor Roberto Roncon-Albuquerque, orientador do projeto, por toda a disponibilidade, pela confiança depositada e pelo incentivo constante, assim como por me ter aberto as portas à atividade científica.

Aos Docentes com quem tive o privilégio de contactar durante o Mestrado Integrado em Medicina, que tanto me inspiraram, nomeadamente ao *Professor Doutor Tiago Henriques-Coelho*, que comigo partilhou o entusiasmo pelos resultados divulgados em abril de 2017, relativos à mais promissora Placenta Artificial apresentada até à data.

Aos Tutores Clínicos que diariamente conjugam a atividade laboral com a vontade de transmitir a sua experiência aos jovens estudantes, por influenciarem e ajudarem a moldar a profissional que espero vir a ser no futuro.

A toda a minha família, em especial aos meus Avós, *Alberto e Maria de Lurdes*, pelo apoio incondicional, pelo carinho e pela tranquilidade que me transmitem a cada dia.

À minha *Mãe*, esperando que esta etapa, que agora termino, possa, de alguma forma, retribuir e compensar toda a dedicação que sempre manifestou para com o meu percurso académico.

Ao meu irmão *Diogo*, companheiro de todas as horas, que celebrou comigo a plenos pulmões cada vitória neste curso e na vida.

Aos meus padrinhos, *Paula e Luís Miguel*, pelo exemplo, e por acompanharem de perto esta longa caminhada.

À *Tatiana* e à *Inês*, por todos os momentos memoráveis dos últimos seis anos, por me terem ajudado a ultrapassar os mais variados desafios e por serem *semper fi*.

Aos *colegas do YES Meeting*, em particular aos “*meus vinte*”, pela honra de poder trabalhar e aprender tanto com eles nessa grande aventura de organizar um congresso, que recordo sempre com o maior dos sorrisos.

À *Margarida* e à *Sofia*, por todo o apoio e amizade, que ultrapassaram largamente a convivência proporcionada pelo YES.

A todos os *demais colegas e amigos* da Faculdade de Medicina e da Faculdade de Engenharia da Universidade do Porto, por terem enriquecido esta jornada.

Ao *Gonçalo*, por acreditar em mim por inteiro e por estar sempre presente.

